



Data reflected in this summary of mosquito-borne disease in the Southwest Region is taken from the Western Australia Notifiable Infectious Disease Database (WANIDD) and includes enhanced surveillance data collected by Population Health Units and Local Governments. (Only locations with notified cases of disease are shown in tables and figures).

Ross River virus (RRV)



There were 9 RRV cases notified by lab during this quarter. These include 5 that were also notified by doctor. No follow data are available. During 2018/19 there were a total of 59 cases which is significantly below the long term monthly mean for all months except August 2018, when it was within the normal range, and June 2019 when it was slightly above the long term monthly mean.

RRV 2019	Apr	May	Jun	Total
SW - Elsewhere	1			1
Donnybrook-Balingup (S)	1			1
DONNYBROOK	1			1
SW - Geographe	1		2	3
Capel (S)	1			1
DALYELLUP	1			1
Busselton (C)			2	2
VASSE			1	1
YALYALUP			1	1
SW - Leschenault		1	4	5
Bunbury (C)			2	2
GLEN IRIS			1	1
WITHERS			1	1
Harvey (S)		1	2	3
AUSTRALIND		1	1	2
COOKERNUP			1	1
Total	2	1	6	9

Barmah Forest Virus (BFV)

There were 2 BFV cases reported by lab during this quarter. One was also notified by doctor. No follow up data are available. During 2018/19 there was a total of 6 cases for this region, this is at or below the long term monthly mean for all months except May 2019 with these two cases. The long term monthly mean is less than two BFV cases per month.


BFV 2019	Apr	May	Jun	Total
SW - Elsewhere		2		2
Manjimup (S)		1		1
WALPOLE		1		1
Nannup (S)		1		1
NANNUP		1		1
Total		2		2






Data reflected in this summary of mosquito-borne disease in the East Metro Region is taken from the Western Australia Notifiable Infectious Disease Database (WANIDD) and includes enhanced surveillance data collected by Population Health Units and Local Governments. (Only locations with notified cases of disease are shown in tables and figures).

Relative Rainfall Mar – May (Autumn) 2019



Relative Rainfall June 2019



Serologically confirmed doctor-notified and laboratory reported cases of Ross River virus disease each month in WA, July 2018 - June 2019

Compiled by the Medical Entomology, WA Department of Health

REGION	Jul	Aug	Sep	Oct	Nov	Dec	Jan	Feb	Mar	Apr	May	Jun	Total	Crude	Age std
KIMBERLEY	3	4	0	0	3	0	1	1	1	1	2	1	17	47.1	58.1
PILBARA	0	1	3	0	0	1	0	0	0	3	3	3	14	22.8	17.2
GASCOYNE	1	0	0	0	0	0	0	0	0	0	0	0	1	10.4	11.2
MIDWEST	0	1	0	0	1	0	0	0	0	1	1	1	5	8.1	7.0
WHEATBELT	2	0	1	1	1	0	1	0	2	0	2	4	14	20.2	18.9
METRO	7	8	12	16	13	7	23	15	13	16	21	11	162	9.4	9.1
PEEL	5	7	13	10	5	8	18	9	9	11	6	5	106	39.8	38.2
LESCHENAULT	1	2	2	3	3	2	3	1	6	0	1	4	28	38.0	36.9
GEOGRAPHE	2	2	3	2	3	2	2	2	4	1	0	2	25	44.6	44.2
ELSEWHERE SW	0	0	0	1	1	0	2	0	1	1	0	0	6	12.7	11.3
SOUTHWEST	8	11	18	16	12	12	25	12	20	13	7	11	165	37.2	
GREAT SOUTHERN	1	1	0	2	0	2	3	1	2	0	1	1	14	23.1	20.9
GOLDFIELDS-ESPERANCE	0	1	2	1	1	2	0	0	2	1	0	1	11	43.2	44.4
WA UNDETERMINED	0	0	0	0	0	0	0	0	0	0	0	0	0		
INTERSTATE	0	0	1	0	2	1	2	2	0	2	0	1	11		
WA TOTAL (does not include interstate)	22	27	36	36	31	24	53	29	40	35	37	33	403		

1) Data current as at 23/07/2019 - table may vary from previous or future versions due to inclusion of additional enhanced surveillance data (Communicable Disease Control Directorate from participating pathology laboratories); Enhanced Surveillance Data (comprising case follow-ups from Environmental Health Officers; patient interviews; Doctor's notifications)

2) Month of onset and suburb/town of exposure determined from Enhanced Surveillance Data where available, and from Doctor's notifications or laboratory reports where not available

3) Data varies from official Western Australian Notifiable Infectious Diseases Database records due to inclusion of Enhanced Surveillance Data (suburb" - (e.g. City of Mandurah unknown)

4) Where a place of exposure occurs in a suburb that carries over 2 Local Governments and it is not clearly defined which local government it occurred in, the case has been entered in the Local Government where the largest portion of the suburb occurs

5) This information is the intellectual property of the Biological and Applied Environmental Health Hazards unit of the WA Department of Health and may not be used for any purpose without prior permission



El Niño conditions are associated with a decrease in rainfall and tidal activity.

La Niña brings wetter and warmer than normal weather which can increase mosquito breeding and mosquito borne diseases.


Positive Indian Ocean Dipole brings below average winter–spring rainfall, above average temperatures.

Australian Bureau of Meteorology (BOM) ENSO issued 9 July 2019

The **El Niño–Southern Oscillation (ENSO)** is currently **neutral** - neither El Niño nor La Niña. While the possibility of El Niño can't be completely ruled out for 2019, the tropical Pacific Ocean is expected to remain in an ENSO-neutral phase over the coming months meaning the **ENSO Outlook remains at INACTIVE**. Model outlooks indicate a **positive Indian Ocean Dipole is likely to be the dominant climate driver for Australia's weather for much of the rest of 2019**, meaning an increased likelihood of a drier than average winter–spring.

International Research Institute for Climate and society IRI ENSO Forecast issued 11 July 2019

The official CPC/IRI outlook, still with an **El Niño advisory**, calls for a 60% chance of ENSO-neutral by Jul-Sep, and neutral remains the most likely category through northern hemisphere fall and winter.



For further information contact

Medical Entomology

PO Box 8172, Perth Business Centre WA 6849

Phone: (08) 9285 5500


Fax: (08) 9383 1819

Email: medical.entomology@health.wa.gov.au




Australian BOM Climate Outlook issued 11 July 2019

A drier than average August to October is likely for large parts of the country, including the northern half of Australia, southwest WA. The month of August is likely to be drier for the far southwest of WA.



Warmer August to October days likely nationwide. Chances are very high (greater than 80%) for the northern half of the country and inland southern WA.



Nights are also likely to be warmer than average for much of Australia. Historical accuracy for August to October maximum temperatures is moderate to high for most of Australia, except for parts of the Pilbara in WA. Minimum temperature accuracy is moderate for most of Australia but patchy across the southeast quarter and western WA.

